e universitaire de santé McGill Il University Health Centre	Patient Information (please print):	MOULET
HME HGM HRV MCH MGH RVH	Date of birth:	MCH File no.
	Last name, First name	
HNM ITM ICL WNH MCI IC		
Montreal Children's Hospital	Current address City, Province	Postal Code
Brain Development Behaviour (BDB) Centralized Intake Referral Form	Home telephone number Other	telephone number
1001, boul. Décarie, Room <b>A04-3140</b> Montreal, Québec, H4A 3J1	Email	
Telephone: (514) <b>412- 4496</b> Fax: (514) <b>412-4136</b> Email:bdbci@muhc.mcgill.ca		
· ·	Language	
Referral date (yyyy/MM/dd):	French ☐ English ☐ Other:	Interpreter needed □
Please describe your concerns:		
Places attach any additional information on a concrete name		
Please attach any additional information on a separate page.		
Please check <u>all</u> that apply:		
☐ Developmental concerns (0-5 years only)		
☐ 1- Child has gross motor and/or fine motor delay	5- Child requires autism evaluation and /o	or autism is suspected for th
<ul><li>□ 2- Child presents with abnormal motor exam (specify belown early handedness</li><li>□ spasticity</li></ul>	w) : following reasons:  ☐ Significant social difficulties	
<ul><li>□ early handedness</li><li>□ spasticity</li><li>□ weakness</li><li>□ weakness with hypotonia</li></ul>	□ Significant social difficulties □ Communication limitations	
☐ early handedness ☐ spasticity ☐ weakness ☐ weakness with hypotonia ☐ other:	☐ Significant social difficulties	
<ul><li>□ early handedness</li><li>□ spasticity</li><li>□ weakness</li><li>□ weakness with hypotonia</li></ul>	☐ Significant social difficulties ☐ Communication limitations ☐ Unusual behaviour / play	of the suspicion of autis
□ early handedness □ spasticity □ weakness □ weakness with hypotonia □ other: □ 3- Child has a speech/language delay □ 4- Child has significant behavioural or emotional difficulties	☐ Significant social difficulties ☐ Communication limitations ☐ Unusual behaviour / play	of the suspicion of autis
□ early handedness □ spasticity □ weakness □ weakness with hypotonia □ other: □ 3- Child has a speech/language delay □ 4- Child has significant behavioural or emotional difficulties Describe: □ □ Hearing test only is required	□ Significant social difficulties □ Communication limitations □ Unusual behaviour / play □ Parents have been informed of	
□ early handedness □ spasticity □ weakness □ weakness with hypotonia □ other: □ 3- Child has a speech/language delay □ 4- Child has significant behavioural or emotional difficulties Describe: □ □ Hearing test only is required	☐ Significant social difficulties ☐ Communication limitations ☐ Unusual behaviour / play	
□ early handedness □ spasticity □ weakness □ weakness with hypotonia □ other: □ 3- Child has a speech/language delay □ 4- Child has significant behavioural or emotional difficulties Describe: □ Hearing test only is required □ Parents suspect hearing loss □ Failed he □ Child presents a high risk of having hearing loss: □ Family history of hearing loss □ Cranio-fa	Significant social difficulties  Communication limitations Unusual behaviour / play  Parents have been informed of the state of the sta	
□ early handedness □ spasticity □ weakness □ weakness with hypotonia □ other: □ 3- Child has a speech/language delay □ 4- Child has significant behavioural or emotional difficulties Describe: □ Hearing test only is required □ Parents suspect hearing loss □ Failed he □ Child presents a high risk of having hearing loss: □ Family history of hearing loss □ Cranio-fa □ Ototoxic medication □ Any med	Significant social difficulties  Communication limitations Unusual behaviour / play  Parents have been informed of the searing test elsewhere. Please specify:  Icial abnormalities  Icial conditions associated with hearing loss	
□ early handedness □ spasticity □ weakness □ weakness with hypotonia □ other: □ 3- Child has a speech/language delay □ 4- Child has significant behavioural or emotional difficulties Describe: □ Hearing test only is required □ Parents suspect hearing loss □ Failed he □ Child presents a high risk of having hearing loss: □ Family history of hearing loss □ Cranio-fa □ Ototoxic medication □ Any medication □ Meningitis Please sp	Significant social difficulties  Communication limitations Unusual behaviour / play  Parents have been informed of the searing test elsewhere. Please specify:  I cial abnormalities ical conditions associated with hearing loss pecify:	
□ early handedness □ spasticity □ weakness □ weakness with hypotonia □ other: □ 3- Child has a speech/language delay □ 4- Child has significant behavioural or emotional difficulties Describe: □ Hearing test only is required □ Parents suspect hearing loss □ Failed he □ Child presents a high risk of having hearing loss: □ Family history of hearing loss □ Cranio-fa □ Ototoxic medication □ Any medi □ Meningitis Please span □ Complicated neonatal course □ Other: □	Significant social difficulties  Communication limitations Unusual behaviour / play  Parents have been informed of the searing test elsewhere. Please specify:  Icial abnormalities  Icial conditions associated with hearing loss	
□ early handedness □ spasticity □ weakness □ weakness with hypotonia □ other: □ 3- Child has a speech/language delay □ 4- Child has significant behavioural or emotional difficulties Describe: □ Hearing test only is required □ Parents suspect hearing loss □ Failed he □ Child presents a high risk of having hearing loss: □ Family history of hearing loss □ Cranio-fa □ Ototoxic medication □ Any medi □ Meningitis □ Please space □ Complicated neonatal course □ Other: □ □ History of otitis media	Significant social difficulties  Communication limitations Unusual behaviour / play  Parents have been informed of the saring test elsewhere. Please specify:  I cial abnormalities iteal conditions associated with hearing loss pecify:	
□ early handedness □ spasticity □ weakness □ weakness with hypotonia □ other: □ 3- Child has a speech/language delay □ 4- Child has significant behavioural or emotional difficulties Describe: □ Parents suspect hearing loss □ Failed he □ Child presents a high risk of having hearing loss: □ Family history of hearing loss □ Cranio-fa □ Ototoxic medication □ Any medi □ Meningitis □ Please sp □ Complicated neonatal course □ Other: □ □ History of otitis media □ Central Auditory Processing evaluation has been recomme	Significant social difficulties  Communication limitations Unusual behaviour / play  Parents have been informed of the saring test elsewhere. Please specify:  I cial abnormalities iteal conditions associated with hearing loss pecify:	
□ early handedness □ spasticity □ weakness □ weakness with hypotonia □ other: □ 3- Child has a speech/language delay □ 4- Child has significant behavioural or emotional difficulties Describe: □ Parents suspect hearing loss □ Failed he □ Child presents a high risk of having hearing loss: □ Family history of hearing loss □ Cranio-fa □ Ototoxic medication □ Any medi □ Meningitis □ Please space □ Complicated neonatal course □ Other: □ □ History of otitis media □ Central Auditory Processing evaluation has been recommed	Significant social difficulties  Communication limitations Unusual behaviour / play  Parents have been informed of the searing test elsewhere. Please specify:  Initial abnormalities ideal conditions associated with hearing loss pecify:  Decided the search of the searc	
□ early handedness □ spasticity □ weakness □ weakness with hypotonia □ other: □ 3- Child has a speech/language delay □ 4- Child has significant behavioural or emotional difficulties Describe: □ Parents suspect hearing loss □ Failed he □ Child presents a high risk of having hearing loss: □ Family history of hearing loss □ Cranio-fa □ Ototoxic medication □ Any medi □ Meningitis □ Please sp □ Complicated neonatal course □ Other: □ □ History of otitis media □ Central Auditory Processing evaluation has been recommed	Significant social difficulties  Communication limitations Unusual behaviour / play  Parents have been informed of the searing test elsewhere. Please specify:  Initial abnormalities ideal conditions associated with hearing loss pecify:  Ended  Family history in a close relative (first decomposition)	egree or sibling of parent)
□ early handedness □ spasticity □ weakness □ weakness with hypotonia □ other: □ 3- Child has a speech/language delay □ 4- Child has significant behavioural or emotional difficulties Describe: □ Parents suspect hearing loss □ Failed he □ Child presents a high risk of having hearing loss: □ Family history of hearing loss □ Cranio-fa □ Ototoxic medication □ Any medi □ Meningitis □ Please sp □ Complicated neonatal course □ Other: □ □ History of otitis media □ Central Auditory Processing evaluation has been recommed □ Risk Indicators □ Child spent time in an NICU after birth □ Child has a microcephaly/macrocephaly	Significant social difficulties  Communication limitations Unusual behaviour / play  Parents have been informed of the searing test elsewhere. Please specify:  Initial abnormalities ideal conditions associated with hearing loss pecify:  Decided the search of the searc	egree or sibling of parent)
□ early handedness □ spasticity □ weakness □ weakness with hypotonia □ other: □ □ 3- Child has a speech/language delay □ 4- Child has significant behavioural or emotional difficulties Describe: □ □ Hearing test only is required □ Parents suspect hearing loss □ Failed he □ Child presents a high risk of having hearing loss: □ Cranio-fa □ Ototoxic medication □ Any medi □ Meningitis □ Please sp □ Complicated neonatal course □ Other: □ □ History of otitis media □ Central Auditory Processing evaluation has been recommed □ Risk Indicators □ Child spent time in an NICU after birth □ Child has a microcephaly/macrocephaly □ Child appears to have dysmorphic features	Significant social difficulties Communication limitations Unusual behaviour / play  Parents have been informed of the parents have been informed have been informed of the parents have been informed have been in	egree or sibling of parent) Il palsy ntal skills are onological age)
□ early handedness □ spasticity □ weakness □ weakness with hypotonia □ other: □ 3- Child has a speech/language delay □ 4- Child has significant behavioural or emotional difficulties Describe: □ Parents suspect hearing loss □ Failed he □ Child presents a high risk of having hearing loss: □ Family history of hearing loss □ Cranio-fa □ Ototoxic medication □ Any medi □ Meningitis □ Please sp □ Complicated neonatal course □ Other: □ □ History of otitis media □ Central Auditory Processing evaluation has been recommed □ Risk Indicators □ Child spent time in an NICU after birth □ Child has a microcephaly/macrocephaly □ Child appears to have dysmorphic features □ Child has experienced an afebrile seizure	Significant social difficulties  Communication limitations Unusual behaviour / play  Parents have been informed of the searing test elsewhere. Please specify:  Initial abnormalities ideal conditions associated with hearing loss pecify:  Ended  Family history in a close relative (first definitellectual disability, autism, or cerebra Child has severe delay (i.e. development)	egree or sibling of parent) Il palsy ntal skills are onological age)
□ early handedness □ spasticity □ weakness □ weakness with hypotonia □ other: □ □ 3- Child has a speech/language delay □ 4- Child has significant behavioural or emotional difficulties Describe: □ □ Hearing test only is required □ Parents suspect hearing loss □ Failed he □ Child presents a high risk of having hearing loss: □ Cranio-fa □ Ototoxic medication □ Any medi □ Meningitis □ Please sp □ Complicated neonatal course □ Other: □ □ History of otitis media □ Central Auditory Processing evaluation has been recommed □ Risk Indicators □ Child spent time in an NICU after birth □ Child has a microcephaly/macrocephaly □ Child appears to have dysmorphic features	Significant social difficulties Communication limitations Unusual behaviour / play  Parents have been informed of the parents have been informed have been informed of the parents have been informed have been in	egree or sibling of parent) Il palsy ntal skills are onological age)